

**Minor impacts of major volcanic eruptions on hurricanes in dynamically-downscaled
last millennium simulations**

Brandon N. Benton,^a Marc J. Alessi,^{a,b} Dimitris A. Herrera,^{a,c} Xiaolu Li,^a Carlos M. Carrillo,^a

Toby R. Ault^a

^a *Cornell University, Ithaca, New York*

^b *Now at Colorado State University, Fort Collins, Colorado*

^c *Now at Instituto Geográfico Universitario, Universidad Autónoma de Santo Domingo,*

Santo Domingo 10103, Dominican Republic

Corresponding author: Brandon Benton, bnb32@cornell.edu; Toby Ault,
toby.ault@cornell.edu

ABSTRACT

The effects of volcanic eruptions on hurricane statistics are examined using two long simulations from the Community Earth System Model (CESM) Last Millennium Ensemble (LME). The first is an unforced control simulation, wherein all boundary conditions were held constant at their 850 CE values (LME_{control}). The second is a “fully forced” simulation with time evolving radiative changes from volcanic, solar, and land use changes from 850 CE through present (LME_{forced}). Large tropical volcanic eruptions produce the greatest change in radiative forcing during this time period, which comprise the focus of this study. The Weather Research and Forecasting (WRF) model is used to dynamically downscale 150 control years of LME_{control} and an additional 84 years of LME_{forced} for all mid-latitude volcanic eruptions between 1100-1850 CE. This time period was selected based on computational considerations. For each eruption, two years are dynamically downscaled. 23 of these volcanic eruptions are in the Northern Hemisphere and 19 are in the Southern Hemisphere. The effectiveness of the downscaling methodology is examined by applying the same downscaling approach to historical ERA-I reanalysis data and comparing the downscaled storm tracks and intensities to the International Best Track Archive for Climate Stewardship (IBTrACS) database. Hurricane statistics are then computed from both the downscaled control and downscaled forced LME simulations. Results suggest moderate effects on hurricanes from the average of all northern hemisphere eruptions, with the largest effects being from the volcanoes with the most aerosol forcing. More specifically, reductions in hurricane frequency, intensity, and lifetime following northern hemisphere eruptions are apparent. Strong evidence is also shown for correlation between eruption strength and changes in these diagnostics. The aggregate effect from both northern and southern hemisphere eruptions is seen to be minor. While reductions in frequency, intensity, and

lifetime from northern hemisphere eruptions occur, the opposite effect is observed from southern hemisphere eruptions.

Keywords: hurricanes, volcanoes, solar geoengineering, last millennium ensemble

1. Introduction

Hurricanes threaten human lives and livelihoods, inflict severe damage to property, and incur billions of dollars in economic losses and recovery efforts. From 1992 through 2011, these events alone caused 42% of the catastrophe-insured losses in the United States (King et al. 2013). The disruption from these extreme weather events will likely increase with rising coastal populations and increasing value of infrastructure in coastal areas (Knutson et al., 2010). Furthermore, anthropogenic climate change is already increasing and expected to further increase average sea surface temperatures (SSTs) and sea level (IPCC, 2016; Kossin et al. 2020). There is also a growing interest in determining if modifications to the incoming flux of solar radiation could potentially offset key impacts expected to occur from rising global temperature (Msadek et al., 2016). Whether or not such strategies are pursued, it is critical to understand the relationship between hurricane statistics and climate responses to past radiative forcings to help characterize the full range of plausible future influences on hurricane activity.

The underlying relationship between hurricanes, radiative forcing, and climate change remains an area of active inquiry (Ting et al., 2015; Elsner, 2006; IPCC, 2014). Several modeling studies have suggested that, in general, future storms may pose more severe threats to human well-being, infrastructure, and the economy (IPCC, 2014). For example, Villarini and Vecchi (2013) and Emanuel (2013) used data from the Coupled Model Intercomparison Project 5 (CMIP5) ensemble to evaluate storm intensity under climate change. Such studies suggest that the number and intensity of the largest storms (e.g., category 4 and 5 hurricanes) will increase in a warmer climate due primarily to an increase in SST. Recent evidence (Kossin et al. 2020) demonstrates that TCs have significantly increased in intensity, though this may be due to changing observational methodologies (Vecchi et al., 2021). Furthermore,

Wang et al. (2018) and Strauss et al. (2021) have shown clear evidence for anthropogenic influence on hurricane precipitation and flooding, respectively.

While we are interested in the relationship between volcanic eruptions and hurricanes for its own sake, eruptions of the recent past and last millennium may also provide a glimpse of the risks associated with “solar geoengineering” (SG), which has received increasing attention recently as a possible strategy for slowing down the rate of global warming from greenhouse gas emissions (Govindasamy and Caldeira, 2000; Caldeira and Wood, 2008; Kravitz et al., 2014; MacMartin et al., 2019). That is, if global greenhouse gas reduction efforts are insufficient in the coming decades, some researchers argue that interventions to the climate system may be preferable to allowing global temperatures to increase (Govindasamy and Caldeira, 2000; Caldeira and Wood, 2008; Kravitz et al., 2014; MacMartin et al., 2019). While there are numerous SG strategies, stratospheric aerosol injection (SAI) in particular mimics large volcanic eruptions (Jones et al., 2017; Crutzen 2006). Fundamentally, SAI decreases the total amount of sunlight reaching the surface, which shares some similarity with the effect of stratospheric aerosol injections from volcanic eruptions. In the stratosphere, chemical and micro-physical processes convert SO_2 into sub-micrometer sulfate particles. The process has been observed in volcanic eruptions (Wilson et al., 1992; Bluth et al., 1992). Although the particle sizes of artificial aerosols are not necessarily the same as volcanic aerosols, the injection rate of artificial aerosols can be designed to roughly match volcanic radiative forcing (Lacis and Mischenko 1995).

Volcanic eruptions result in an increased reflection of solar radiation, which can strongly impact global temperatures, circulation patterns, and water cycles (Mass and Portman, 1989; Robock, 2000; Swingedouw et al., 2017). Changes in tropical sea surface temperature from SAI and volcanic eruptions would also similarly alter the position of the

Inter Tropical Convergence Zone (ITCZ). For example, asymmetric volcanic forcings (e.g., from volcanic aerosols being concentrated in the stratosphere of one hemisphere) would alter the position of the ITCZ for at least one year following an eruption (Jones et al., 2017). These effects could potentially be even longer lasting if coupled interactions between the ocean and atmosphere are engaged (Colose et al., 2016; Raible et al., 2016; Stevenson et al., 2016; Schurer et al., 2014; Schurer et al., 2013). Given that hurricanes are sensitive to the regions where moisture convergence occurs, it follows that such radiative effects on the global circulation could influence hurricane statistics.

Comparisons between all-forcing and single-forcing last millennium model simulations, as well as multiproxy paleoclimate constructions, have shown that large volcanic eruptions were the dominant forcing during the pre-industrial record (Schurer et al., 2014; Schurer et al., 2013). Studies have also shown that large tropical volcanic eruptions may have long-lasting influences on the Atlantic multi-decadal oscillation and lead to El Niño-like warming in the cold season after the eruption (Stevenson et al., 2016; Otto-Bliesner et al., 2016). Such impacts could, in turn, affect hurricane statistics because these large-scale modes help govern, in part, the frequency and intensity of storms occurring in any given year.

The historical record only provides a few clues about the effect of volcanic eruptions on hurricanes or tropical cyclones (TCs). Nevertheless, modeling studies suggest that a reduction in TC accumulated energy, TC duration, and lifetime maximum intensity occurs following a volcanic eruption due to a decrease in SST and increase in upper tropospheric/lower stratospheric temperature (Evan, 2012), all of which decreases TC thermodynamic efficiency estimates (Emanuel et al., 2013). There is evidence that after some eruptions, an asymmetric increase in stratospheric aerosols occurs in the hemisphere in which the eruption took place, modifying the sea surface temperature gradient (Haywood et al.,

2013). This gradient shifts the location of the Inter-tropical Convergence Zone (ITCZ) to the opposite hemisphere of the eruption (Haywood et al., 2013), which hinders hurricane development in the volcano's own hemisphere due to a decrease in convection and increase in wind shear. In fact, after the northern hemisphere eruptions of Mount Pinatubo (1991) and El Chichón (1982), North Atlantic TC activity decreased, while TC activity increased following the southern hemisphere eruption of Agung (1964) (Evan, 2012; Guevara-Murua et al., 2015). More recently, Camargo and Polvani (2019) found no robust reduction of North Atlantic tropical cyclone activity in recent observations or reanalysis. A reduction in potential intensity (PI) is observed in CMIP5 and CESM large-ensemble historical simulations after volcanic eruptions, though the decrease might be overestimated due to model bias.

Attempts to use paleoclimate indicators of past events (i.e., paleotempestology) are limited by the paucity of appropriate archives and proxies (Liu et al., 2008; Mann et al., 2009; Donnelly et al., 2015). Importantly, most paleorecords are commonly constrained to certain geographic areas and currently provide limited information from other regions also commonly affected by TCs, such as the Caribbean (Oliva et al., 2018). The use of paleotempestological records is further limited because it is not possible to fully reconstruct the tracks and lifetime of past TCs, most of which occur over the ocean (Emanuel, 2005). These limitations preclude the extensive use of such data to evaluate the effects of major volcanic eruptions on TC activity (Oliva et al., 2018; Yan et al., 2015; Korty et al., 2012).

Hurricanes are mesoscale features of the tropical circulation, and as such they depend critically on quantities that are typically unresolvable in the coarse resolution grid of general circulation models (GCMs), which typically have nominal horizontal resolutions on the order of 50-200km. For example, the resolution of the CESM simulation used here is $1.9^{\circ} \times 2.5^{\circ}$ which is roughly 211km by 278km at the equator. In order to overcome this limitation,

authors typically follow one of three following approaches. The first approach is relatively simple, and entails calculating thermodynamic metrics like PI at the native (coarse) resolution of GCM output, which shows the maximum possible TC intensity given a vertical sounding (Wang et al., 2018; Emanuel and Nolan, 2004; Tang and Camargo, 2014; Bister and Emanuel, 2002). A recent study (Yan et al., 2018) used the last millennium ensemble (LME) to determine the theoretical effects of volcanic eruptions during the last 1000 years on PI. The authors found a significant relationship lasting up to 3 years post-eruption, but also “divergent” responses at the mid and high latitudes to the volcanic forcing. While this approach is computationally efficient, it does not explicitly simulate TCs.

A second approach uses a statistical method, coupling synthetic tracks with an axisymmetric hurricane model to downscale GCM output (Emanuel, 2006; Kerty et al., 2017). This approach is computationally lightweight, allowing for an investigation of long-term variability in fully coupled GCM simulations of the last millennium (Kozar et al., 2013). While this method has a high spatial resolution in the vicinity of the eyewall, the model is limited due to being axisymmetric and a hurricane being unable to feedback on environmental conditions supplied to it.

The third approach employs a regional model to dynamically downscale GCM output (Knutson et al., 2008). Dynamical downscaling typically requires high performance computing infrastructure as well as boundary conditions from the “parent” GCM at six hourly temporal resolution. It is therefore much more computationally expensive than the other two methods, but it provides greater insight into the storms that would have occurred in each GCM framework if it were run with sufficiently high spatial resolution. Dynamical downscaling has been widely used to evaluate hurricane statistics during the 20th and 21st

century (Emanuel, 2013), but it has not been widely adapted to the last millennium modeling context.

In this study, we use WRF to dynamically downscale output from the LME to analyze the effect of volcanic eruptions on hurricane intensity and lifetime. While not a perfect analog to SAI, eruptions are important for understanding physical mechanisms and validating models used in projecting the climate response to a possible SAI regime. Understanding the effects of volcanic eruptions on hurricane dynamics during the last millennium can therefore yield some insight into the possible (unintended) effects of SAI on these hazards.

2. Data and Methods

We dynamically downscale 234 total years from two members of the “Last Millennium Ensemble” (Otto-Bliesner et al., 2016) with high resolution temporal output. The LME consists of over two dozen fully forced and single forcing experiments from the record spanning 850 CE to 2005. The LME was used because it provided an extensive readily available dataset which could be easily downscaled. While monthly data was archived for most of the members of the LME, only two simulations produced sufficiently high temporal output to allow for high resolution dynamical downscaling using a regional model. One of these runs was a fully forced last millennium simulation and the other was a long control simulation with time invariant boundary conditions. We further evaluate the strengths and limitations of our methodology by comparing downscaled reanalysis data to an historical database of hurricane tracks and intensities. The details of this approach are described below.

a. Data

1) Last Millennium Ensemble

Output from only two members of the LME were archived at sufficiently high temporal resolution to be used as boundary conditions for WRF: a fully-forced simulation with time varying boundary conditions (LME_{forced}) and a pre-industrial control simulation with time invariant boundary conditions (LME_{control}). Both simulations were run from 850 to 2005 CE using the Community Earth System Model (CESM) version 1.1, with the Community Atmosphere Model (CAM) version 5. The resolution of the atmosphere and land grids are nominally $\sim 2^\circ$, and $\sim 1^\circ$ for ocean and sea ice grids. While both runs were spun up for 200 years under control conditions prior to 850 CE, LME_{forced} was forced with the transient evolution of solar intensity, volcanic emissions, greenhouse gases, aerosols, and land-use conditions, as well as insolation changes from planetary orbit and tilt. In the LME_{control} the boundary conditions were simply held fixed at their pre-industrial levels, thus providing an unforced baseline for evaluating changes in hurricane statistics following large volcanic eruptions.

2) ERA-I and IBTrACS

The native 2° resolution of CAM5 in the LME simulations would make it impossible to resolve hurricanes, hence we cannot evaluate the reliability of our downscaling methodology (see section 2.b.1) with LME data alone. We therefore also downscaled the ERA-Interim (ERA-I) (Dee et al., 2011) reanalysis data to retrospectively predict historical hurricanes, then compared those predictions against the International Best Track Archive for Climate Stewardship (IBTrACS) (Knapp et al., 2010) database. ERA-I comprises a reanalysis dataset starting in 1979 and is available until August 2019. It uses four-dimensional variational data assimilation (4DVAR), yielding a significant advantage over reanalysis products using 3DVAR. This improves synoptic data handling and allows for the influence of an observation to be more strongly controlled by model dynamics (Schenkel and Hart,

2011). This data assimilation method is coupled with the ECMWF Integrated Forecast Model (IFS) to extrapolate fields between observations.

b. Methods

1) Dynamical Downscaling

We used WRF version 3.9 (WRFV3.9) (Skamarock et al., 2008) to dynamically downscale archived data from LME simulations with the physics schemes shown in Table 1. We also turned on heat and moisture surfaces fluxes ($isflx=1$) and modified exchange coefficients C_d and C_k according to surface winds ($isftcflx=1$). The physics schemes were selected to satisfy a 15% difference threshold imposed between downscaled ERA-I and IBTrACS, as quantified by our diagnostics suite.

WRF was run for a total of 234 simulation years over a domain spanning the North American sector from 130°W to 15°E and the equator to 55°N , allowing us to identify and track TCs in both the Atlantic and Eastern Pacific, even after making landfall in North America. In comparing LME_{control} and LME_{forced} we focused on the effect of aerosol forcing from volcanic eruptions. The aerosol mass signals from volcanic eruptions from 1100-1850 CE are shown in Figure 1, which is described in detail in Gao et al. (2008). We selected the top 42 eruptions based on peak aerosol mass signal from LME_{forced} and ran WRF for two years after each of these eruptions (84 simulation years total). LME_{control} was run using WRF for 150 years from 1000 to 1150 to provide a sufficient sample of non-volcanic natural variability. The two year window after eruptions was selected because the residence time of stratospheric aerosols is around one to two years (Crutzen 2006), and individual large eruptions produce global or hemispheric cooling for an average of two or three years (Robock 2015). The LME_{control} run was ensured to have sufficient length by analyzing the SST signal

in frequency space (Figure 2). This analysis shows that using 100 years of control data is sufficient to ensure we are not missing any low frequency variability.

We elected to use a horizontal grid spacing (ΔX) of 30km. The 30km spacing represents a compromise between our competing requirements for high resolution output and a large sample size; each downscaled year used approximately 2,000 core hours on the Cheyenne supercomputer totaling about 500,000 core hours for all years. While the 30km resolution is somewhat coarse for resolving certain features of hurricanes, decreasing the resolution further to 10km was much too expensive. The threefold increase in spatial resolution would have translated into more than a 10-fold increase in core hours, or a 10-fold reduction in the number of years simulated. All data from the LME were prepared for WRF using procedure and code described in (Bruyere et al., 2015).

In our work, 6- hourly ERA-I data was also downscaled in WRF (using the same domain and 30km resolution as the downscaled LME simulations) from 1995 to 2005 and compared to the IBTrACS database for the same record. The comparison was made using the suite of diagnostics described in section 2.b.3.

2) Tracking Tropical Cyclones with TSTORMS

We applied the TSTORMS (Zhao et al., 2009) tracking software, developed and supported by GFDL, to analyze the results of downscaling. This routine uses minimum pressure and maximum vorticity criteria to identify cyclones. Events are stored as “storms” if they satisfy the following conditions for at least a preset number of days (ndays): (1) That the maximum vorticity location is within a threshold radius (rcrit) of the minimum pressure location, (2) that the core temperature of the cyclone is higher than outside of the core by a threshold difference (twccrit) and (3) the difference in vertical distance between pressure

levels at 200hPa and 1000hPa outside and inside the core exceeds a threshold value (thickcrit).

As described in Walsh et al. (2015) and Zhao et al. (2009), tracking results are sensitive to the details of the tracking scheme that is employed and especially the threshold values selected for identifying storms (Horn et al., 2014). To identify sensitivity to threshold values, we conducted a limited parameter sweep to determine optimal threshold values. We calculated the difference between ERA-I downscaled output and IBTrACS data, for each set of parameters, using the diagnostics described in section 2.b.3. We used the set of parameters that achieved the minimum difference of $\sim 13.5\%$. This parameter set was $rcrit = 1.5^\circ$, $twccrit = 1.0^\circ\text{C}$, $thickcrit = 50\text{m}$, and $ndays = 2$.

3. Diagnostics

Once hurricanes were identified in our downscaled LME data using TSTORMS, we calculated 15 diagnostic metrics to evaluate differences in the statistics of storms in LME_{control} and those occurring after large eruptions in LME_{forced} . These diagnostics consist of fraction of storms vs (1) month, (2) year, (3) latitude, (4) longitude, (5) maximum wind speed, (6) minimum pressure, (7) decay time from maximum wind speed, (8) decay time from minimum pressure, (9) power dissipation index (PDI), and (10) accumulated cyclone energy (ACE). For example, (5) is a plot of the fraction of storms with a specific maximum wind speed. Additionally, we calculated percentage of storms within (10) May to November, (11) $0 - 25\text{N}$ latitude, (12) $100\text{W} - 50\text{W}$ longitude, (13) $1020\text{hPa} - 980\text{hPa}$ pressure, (14) $0\text{m/s} - 40\text{m/s}$ maximum wind speed, (15) $0 - 100\text{hrs}$ decay time from maximum wind speed, (16) $0 - 100\text{hrs}$ decay time from minimum pressure, (17) $1.12 \cdot 10^9 - 21.6 \cdot 10^9 \text{m}^3/\text{s}^2$ PDI, and (18) $0.25 - 3.75 \text{m}^2/\text{s}^2$ ACE. Mean values and quantile values (expressed as percentages) were used to calculate fractional differences between LME_{control} and LME_{forced} , and these differences

were averaged over all diagnostics for a composite percentage difference. We refer to the mean difference of diagnostics 1-8 as the total “average difference” and the mean difference of diagnostics 9-15 as the total “percentage difference.” We also computed these metrics from our downscaled ERA-I data to compare them to the IBTrACS database as a test of our methodology, as described in sections 2.b.1 and 2.b.2.

The diagnostics described above were used as test statistics to evaluate whether volcanic eruptions have a measurable effect on hurricane behavior. These diagnostics were selected in order to assess hurricane behavior across a broad range of characteristics. The diagnostics not only quantify hurricane behavior across the temporal and spatial domain, but also assess more fundamental physical characteristics. In addition, the diagnostics can be used with limited data consisting only of time, location, wind speed, and surface pressure. This presents a versatile and efficient approach to capture both mean climatology and structured hurricane behavior.

To determine whether volcanic eruptions affect hurricane statistics, we performed two-sample KS-tests for distributions of each of the diagnostics. The two samples tested for each diagnostic came from downscaled LME_{control} and LME_{forced} data. Since LME_{control} does not include volcanic eruptions, agreement with LME_{control} is confirmation of the null hypothesis. We also performed two-sample Anderson-Darling tests to account for long tail effects to which KS-tests are insensitive.

3. Results

We first show comparisons in TC tracks between downscaled output from ERA-I and IBTrACS (Figure 3) as well as between LME_{forced} and LME_{control} (Figure 4). The ERA-I vs. IBTrACS comparison provides a baseline of whether WRF downscales TCs correctly, while

the comparison between LME_{forced} and LME_{control} is focused specifically on the effect of aerosol forcing from volcanic eruptions. There is agreement in the location of TC tracks for both comparisons, however the downscaled ERA-I underestimates TC intensity. This underestimation is likely related to the downscaling resolution of 30km and the representation of storms in ERA-I (Lui et al. 2021). There is also a notable lack of TC tracks in the Gulf of Mexico for the downscaled ERA-I compared to the IBTrACS.

a. ERA-I vs. IBTrACS

As shown in Table 2, using our suite of diagnostics, we found an overall agreement between ERA-I and IBTrACS of $\sim 86.5\%$, or a composite difference of $\sim 13.5\%$. 6-hourly ERA-I data downscaled in WRF was compared to IBTrACS for the same time period (1995 – 2005). Notably, most hurricanes (97%) from downscaled ERA-I remain below a maximum wind speed of 40 m/s, whereas 30% of storms from IBTrACS cross the 40 m/s maximum wind speed. Diagnostic distributions for both ERA-I and IBTrACS are shown in Figure 6. It is worth noting that the truncation of the domain in our ERA-I WRF simulations contributes to the differences in latitude and longitude peaks seen in Figure 6. For example, according to the IBTrACS database, some storms track beyond the WRF domain, while our downscaled ERA-I simulations cut-off where the WRF domain ends. In Hodges et al. (2017), the authors assess how well TCs are represented in reanalysis products. This work used two TC-track matching approaches, referred to as (1) “direct matching” and (2) “objective matching”. The authors further used several diagnostics in order to compare reanalysis TC tracks to those found in IBTrACS. The objective matching approach, which employs a tracking algorithm similar to TSTORMS, found an agreement of $\sim 60\%$ with ERA-I in the Northern Hemisphere. A simple “direct matching” implementation of our own achieved similar agreement. The physics schemes in section 2.b.1 and threshold values in section 2.b.2 were used to satisfy a

self-selected 15% difference threshold imposed between ERA-I and IBTrACS, as quantified by our diagnostics suite.

b. Effect of Eruptions on Hurricane Statistics

1) Average Effect of Eruptions

Distributions of the diagnostics for LME_{control} and LME_{forced} , with all 42 eruptions included, are shown in Figure 7. The same diagnostic comparison for all northern hemisphere eruptions is shown in Figure 8. All southern hemisphere eruptions are included in the diagnostic comparison shown in Figure 9. A diagnostic comparison for the top 5 northern hemisphere eruptions is shown in Figure 10. The top 5 southern hemisphere eruptions are included in Figure 11. Performing two sample KS-tests and two-sample Anderson-Darling tests on the distributions, along with fractional significance tests on the difference of mean values, shows that the overall effect of all 42 eruptions is consistent with the null hypothesis. That is, the effect of all 42 eruptions is consistent with the natural climate variability seen in LME_{control} (Table 3). However, when separating northern and southern hemisphere eruptions, we observe a moderate effect on the frequency, lifetime, and intensity of hurricanes (Tables 4, 6). The KS-tests show a maximum difference between the two samples (D-value) and a probability that the two samples are drawn from the same distribution (P-value). The Anderson-Darling tests give the level at which the null hypothesis can be rejected. For example, in the max wind row of Table 6, an Anderson-Darling test significance value of 0.001 shows that the null hypothesis can be rejected at the 0.1% level. For all northern hemisphere eruptions (Table 4) we can reject the null hypothesis by the Anderson-Darling test for intensity, PDI, and ACE at a maximum level of 13%. For both PDI and ACE we can reject the null hypothesis at around the 2% level. We see similar results for all southern

hemisphere eruptions in Table 6. The fractional significance tests (Tables 8-10) show the fraction of TCs in LME_{control} which exceed the LME_{forced} mean. For example, in the max wind row of Table 8, 53.2% of the TCs in LME_{control} had stronger maximum wind speed than the mean maximum wind speed of TCs in LME_{forced} . For all northern hemisphere eruptions, we see in Table 9 that TCs have a lower lifetime, maximum intensity, and latitude than 60-70% of TCs in LME_{control} . We also find that TCs following northern hemisphere eruptions have a lower PDI and ACE than 65% of TCs in the control. We observe the opposite effect for southern hemisphere eruptions. We see in Table 10 that nearly 60% of TCs in the control have longer lifetimes, larger maximum intensity, increased latitudes, and larger PDI/ACE than TCs following southern hemisphere eruptions.

We also calculated pearson correlation coefficients on eruption strength and diagnostic changes. To determine the significance of these correlation coefficients we also calculated the 90%, 85%, and 80% confidence intervals. If the confidence interval for a particular diagnostic (e.g. maximum wind speed) does not include zero correlation we can say with at least 90%, 85%, or 80% confidence that there is a correlation between eruption strength and that particular diagnostic. These confidence intervals are affected by the number of eruptions in our analysis. The correlation coefficients for all eruptions, in both hemispheres, are listed in Table 11. Here we see with at least 80% confidence that there is a correlation between eruption strength and reduction in yearly hurricane number, intensity, and lifetime. The correlation coefficients for all northern hemisphere eruptions are shown in Table 12. Again we see with at least 80% confidence that there is a correlation between eruption strength and reduction in yearly hurricane number, intensity, and lifetime. The correlation coefficients for all southern hemisphere eruptions are shown in Table 13. Here we again see with at least 80% confidence there is a correlation between eruption strength and reduction in

intensity. We also see with at least 90% confidence a correlation between eruption strength and storms occurring earlier in the year. However, for the southern hemisphere we do not see the same correlation between eruption strength and reduction in yearly number or lifetime. For the southern hemisphere we see with at least 90% confidence a correlation between eruption strength and storm occurring later in the year. The opposite effect is seen for northern hemisphere eruptions.

2) Effect of Strongest Eruptions

Distributions of diagnostics with only the 5 strongest northern hemisphere eruptions are shown in Figure 10. The same diagnostics for the 5 strongest southern hemisphere eruptions are shown in Figure 11. The results of running KS-tests and Anderson-Darling tests for the 5 strongest northern hemisphere eruptions are shown in Table 5. The results for the 5 strongest southern hemisphere eruptions are shown in Table 7. For the 5 largest eruptions, in both the northern and southern hemispheres we can reject the null hypothesis by the Anderson-Darling test for nearly every diagnostic at a maximum level of 1%. The fractional significance tests on the strongest northern hemisphere and southern hemisphere eruptions are in Tables 9 and 10, respectively. We see in Table 9 that for the 5 strongest northern hemisphere eruptions 70-80% of TCs have a lower maximum intensity, shorter lifetime, and are found at a lower latitude than TCs in LME_{control} . Following the strongest eruption (1258) nearly 80-90% of TCs exhibit this reduced lifetime, maximum intensity, and latitude. We see in Table 10 that for the 5 strongest southern hemisphere eruptions there is the opposite effect to that of northern hemisphere eruptions. In this case nearly 60% of TCs following eruptions have an increased lifetime, maximum intensity, and latitude. However, we still observe a similar reduction in power dissipation index and accumulated cyclone energy.

4. Discussion and Conclusion

In this work we have explored the effect of volcanic eruptions in the past millennium on hurricane climatology. To do this we first validated our approach of downscaling CESM data with WRF by comparing results of ERA-I downscaling with IBTrACS data. ERA-I downscaling was only able to partially reproduce IBTrACS observations however, underestimating observational intensities. On the other hand, frequency, lifetime, and location were faithfully reproduced in the ERA-I downscaling. The underestimation of intensity could limit measurable effects in our intensity-based diagnostics. We also performed a parameter search for our cyclone tracking algorithm in order to achieve high accuracy. We then compared the results of downscaling our control data from CESM with forced data from CESM, where we focused on the years in the forced data with volcanic eruptions.

Our results suggest that the overall effect of *all* eruptions on hurricane statistics is small and not significant as compared to the control simulation (i.e., the null hypothesis). However, we see a moderate reduction in frequency, lifetime, and maximum intensity for hurricanes following northern hemisphere eruptions and the opposite for southern hemisphere eruptions. Sufficiently strong northern hemisphere eruptions do result in lower annual hurricane count, reduced intensity, and shorter lifetimes, significant at the 70-80th percentile. This evidence is in the form of KS, Anderson-Darling, and fractional significance tests on diagnostic distributions, as well as correlations between strength and changes in the mean values of these diagnostics. The moderate increase in North Atlantic TC activity following southern hemisphere eruptions substantiates previous research suggesting moisture convergence increases in the hemisphere opposite of a volcanic eruption (e.g. Haywood et al., 2013).

Comparing the downscaled ERA-I results to hurricane tracks and intensities from IBTrACS allowed us to evaluate the efficacy of our approach. Due to the inherently chaotic nature of hurricane genesis, exact agreement between ERA-I and IBTrACS was not expected, especially given the coarse 30km resolution of our downscaled simulations. Assessing our approach was the primary objective in comparing ERA-I with IBTrACS. We expected ERA-I to capture the observational record for mean climate and to provide good agreement between downscaled results and overall hurricane statistics seen in IBTrACS. We avoided using metrics like PI, since caution is urged in using PI to draw strong conclusions about tropical cyclone projections as it fails to capture features seen in high-resolution climate models (Wehner et al., 2015). Dynamical downscaling provides far greater detail in both the spatial and temporal domain (Emanuel, 2013). This is evident in Figures 4 and 5 and the extensive suite of diagnostics used to analyze downscaled results.

The results in this study have significant implications for hurricane development in a potential future climate under an SAI regime. Although we analyzed the effects of an increase in stratospheric aerosols from volcanic eruptions, the results are relevant to what may occur under an SAI regime. Although analyses of impacts were once limited by historical observation and coarser resolution, we were able to evaluate the direct influence of many volcanic eruptions on individual hurricanes. For example, northern hemisphere eruptions in the downscaled LME_{forced} experiment produced a slight reduction in hurricane frequency, intensity, and lifetime. These impacts could be similarly felt if SAI was implemented, removing some uncertainty associated with regional changes in tropical cyclone development for the Northern Atlantic Ocean. Under a relatively strong SAI regime and according to our results, hurricanes would likely decrease in frequency, lifetime, and intensity.

Although our results show moderate correlation between eruption strength and certain diagnostic measures, it is not necessarily true that stronger eruptions have a larger effect on hurricane statistics. Additionally, research has shown large uncertainties in volcanic reconstructions and seasonality of volcanic eruptions (Schmidt et al., 2012; Stevenson et al., 2017; Raible et al., 2016). This presents a direction for further investigation. In this vein, one could look at an ensemble of higher resolution GCM simulations on one or two of the strongest volcanic eruptions. This eruption profile could be simulated both in the climate conditions during the historical eruption as well as under future climate change conditions. An ensemble average of simulations with perturbed initial conditions will allow us to home in on the sole effect of aerosol forcing. This will also allow us to explore the question of whether downscaling introduced any unknown biases. An ensemble under future climate change conditions will allow us to explore the interplay of large aerosol forcing and strong anthropogenic forcing.

Acknowledgments.

We are grateful for the early assistance we received from Dr. Bette Otto-Bliesner in setting up the original LME control as well as to Yu Izuka for his initial work on this project. Additionally we thank the reviewers for numerous helpful comments which significantly improved the paper. We would like to acknowledge high-performance computing support from Cheyenne ([doi:10.5065/D6RX99HX](https://doi.org/10.5065/D6RX99HX)) provided by NCAR's Computational and Information Systems Laboratory, sponsored by the National Science Foundation. Both Dr. Benton and Dr. Ault were partially supported by NSF Grants AGS 1602564 and AGS 1751535. The authors declare no conflicts of interest with this publication.

REFERENCES

- Bister, M., & Emanuel, K. A. (2002). Low frequency variability of tropical cyclone potential intensity 1. Interannual to interdecadal variability. *Journal of Geophysical Research: Atmospheres*, 107(D24), ACL 26-1-ACL 26-15.
<https://doi.org/https://doi.org/10.1029/2001JD000776>
- Bluth, G. J. S., Doiron, S. D., Schnetzler, C. C., Krueger, A. J., and Walter L. S. (1992), Global tracking of the SO₂ clouds from the June 1991 Mount Pinatubo eruptions, *Geophysical Research Letters* 19, 151–154
- Bruyere, C. L., Monaghan, A. J., Steinhoff, D. F., & Yates, D. (2015). *Bias-Corrected CMIP5 CESM Data in WRF/MPAS Intermediate File Format*.
<https://doi.org/10.5065/D6445JJ7>
- Caldeira, K., & Wood, L. (2008). Global and Arctic climate engineering: numerical model studies. *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 366(1882), 4039–4056.
<https://doi.org/10.1098/rsta.2008.0132>
- Camargo, S. J., & Polvani, L. M. (2019). Little evidence of reduced global tropical cyclone activity following recent volcanic eruptions. *Npj Climate and Atmospheric Science*, 2(1), 14. <https://doi.org/10.1038/s41612-019-0070-z>

- Colose, C. M., LeGrande, A. N., & Vuille, M. (2016). Hemispherically asymmetric volcanic forcing of tropical hydroclimate during the last millennium. *Earth System Dynamics*, 7(3), 681–696. <https://doi.org/10.5194/esd-7-681-2016>
- Crutzen, P.J. (2006). Albedo Enhancement by Stratospheric Sulfur Injections: A Contribution to Resolve a Policy Dilemma?. *Climatic Change* 77, 211.
<https://doi.org/10.1007/s10584-006-9101-y>
- Dee, D. P., Uppala, S. M., Simmons, A. J., Berrisford, P., Poli, P., Kobayashi, S., Andrae, U., Balmaseda, M. A., Balsamo, G., Bauer, P., Bechtold, P., Beljaars, A. C. M., van de Berg, L., Bidlot, J., Bormann, N., Delsol, C., Dragani, R., Fuentes, M., Geer, A. J., ... Vitart, F. (2011). The ERA-Interim reanalysis: configuration and performance of the data assimilation system. *Quarterly Journal of the Royal Meteorological Society*, 137(656), 553–597.
<https://doi.org/https://doi.org/10.1002/qj.828>
- Donnelly, J. P., Hawkes, A. D., Lane, P., MacDonald, D., Shuman, B. N., Toomey, M. R., van Hengstum, P. J., & Woodruff, J. D. (2015). Climate forcing of unprecedented intense-hurricane activity in the last 2000 years. *Earth's Future*, 3(2), 49–65. <https://doi.org/https://doi.org/10.1002/2014EF000274>
- Dyer, A. J., & Hicks, B. B. (1970). Flux-gradient relationships in the constant flux layer. *Quarterly Journal of the Royal Meteorological Society*, 96(410), 715–721.
<https://doi.org/https://doi.org/10.1002/qj.49709641012>

- Elsner, J. B. (2006). Evidence in support of the climate change–Atlantic hurricane hypothesis. *Geophysical Research Letters*, 33(16).
<https://doi.org/https://doi.org/10.1029/2006GL026869>
- Emanuel, K. A. (1995). On Thermally Direct Circulations in Moist Atmospheres. *Journal of Atmospheric Sciences*, 52(9), 1529–1534.
[https://doi.org/10.1175/1520-0469\(1995\)052<1529:OTDCIM>2.0.CO;2](https://doi.org/10.1175/1520-0469(1995)052<1529:OTDCIM>2.0.CO;2)
- Emanuel, K. A. (2005). Increasing destructiveness of tropical cyclones over the past 30 years. *Nature*, 436(7051), 686–688. <https://doi.org/10.1038/nature03906>
- Emanuel, K. A. (2006). Climate and Tropical Cyclone Activity: A New Model Downscaling Approach. *Journal of Climate*, 19(19), 4797–4802.
<https://doi.org/10.1175/JCLI3908.1>
- Emanuel, K. A. (2013). Downscaling CMIP5 climate models shows increased tropical cyclone activity over the 21st century. *Proceedings of the National Academy of Sciences*, 110(30), 12219 LP – 12224. <https://doi.org/10.1073/pnas.1301293110>
- Emanuel, K. A., & Nolan, D. S. (2004). Tropical cyclone activity and the global climate system. *26th Conference on Hurricanes and Tropical Meteorology*.
- Emanuel, K., Solomon, S., Folini, D., Davis, S., & Cagnazzo, C. (2013). Influence of Tropical Tropopause Layer Cooling on Atlantic Hurricane Activity. *Journal of Climate*, 26(7), 2288–2301. <https://doi.org/10.1175/JCLI-D-12-00242.1>

- Evan, A. T. (2012). Atlantic hurricane activity following two major volcanic eruptions. *Journal of Geophysical Research: Atmospheres*, *117*(D6).
<https://doi.org/https://doi.org/10.1029/2011JD016716>
- Gao, C., Robock, A., & Ammann, C. (2008). Volcanic forcing of climate over the past 1500 years: An improved ice core-based index for climate models. *Journal of Geophysical Research: Atmospheres*, *113*(D23).
<https://doi.org/https://doi.org/10.1029/2008JD010239>
- Govindasamy, B., & Caldeira, K. (2000). Geoengineering Earth's radiation balance to mitigate CO₂-induced climate change. *Geophysical Research Letters*, *27*(14), 2141–2144. <https://doi.org/https://doi.org/10.1029/1999GL006086>
- Guevara-Murua, A., Hendy, E. J., Rust, A. C., & Cashman, K. V. (2015). Consistent decrease in North Atlantic Tropical Cyclone frequency following major volcanic eruptions in the last three centuries. *Geophysical Research Letters*, *42*(21), 9425–9432. <https://doi.org/https://doi.org/10.1002/2015GL066154>
- Haywood, J. M., Jones, A., Bellouin, N., & Stephenson, D. (2013). Asymmetric forcing from stratospheric aerosols impacts Sahelian rainfall. *Nature Climate Change*, *3*(7), 660–665. <https://doi.org/10.1038/nclimate1857>
- Hodges, K., Cobb, A., & Vidale, P. L. (2017). How Well Are Tropical Cyclones Represented in Reanalysis Datasets? *Journal of Climate*, *30*(14), 5243–5264.
<https://doi.org/10.1175/JCLI-D-16-0557.1>

- Hong, S., & Lim, J. (2006). The WRF Single-Moment 6-Class Microphysics Scheme (WSM6). *Asia-Pacific Journal of Atmospheric Sciences*, 42, 129–151.
- Hong, S.-Y., Noh, Y., & Dudhia, J. (2006). A New Vertical Diffusion Package with an Explicit Treatment of Entrainment Processes. *Monthly Weather Review*, 134(9), 2318–2341. <https://doi.org/10.1175/MWR3199.1>
- Horn, M., Walsh, K., Zhao, M., Camargo, S. J., Scoccimarro, E., Murakami, H., Wang, H., Ballinger, A., Kumar, A., Shaevitz, D. A., Jonas, J. A., & Oouchi, K. (2014). Tracking Scheme Dependence of Simulated Tropical Cyclone Response to Idealized Climate Simulations. *Journal of Climate*, 27(24), 9197–9213. <http://www.jstor.org/stable/26194640>
- Iacono, M. J., Delamere, J. S., Mlawer, E. J., Shephard, M. W., Clough, S. A., & Collins, W. D. (2008). Radiative forcing by long-lived greenhouse gases: Calculations with the AER radiative transfer models. *Journal of Geophysical Research: Atmospheres*, 113(D13). <https://doi.org/https://doi.org/10.1029/2008JD009944>
- IPCC. (2018). *Global Warming of 1.5 degrees Celsius: An IPCC Special Report on the Impacts of Global Warming of 1.5 degrees C Above Pre-industrial Levels and Related Global Greenhouse Gas Emission Pathways, in the Context of Strengthening the Global Response to the Thr.*
- IPCC. (2014). *Climate Change 2014: Synthesis Report. Contribution of Working Groups I, II and III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change.* <https://www.ipcc.ch/report/ar5/syr/>

IPCC. (2007). *Climate Change 2007: Synthesis Report. Contribution of Working Groups I, II and III to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change.*

Jones, A.C., Hawcroft, M.K., Haywood, J.M., Jones, A., Guo, X. and Moore, J.C. (2018), Regional Climate Impacts of Stabilizing Global Warming at 1.5 K Using Solar Geoengineering. *Earth's Future*, 6: 230-251.
<https://doi.org/10.1002/2017EF000720>

Jones, A. C., Haywood, J. M., Dunstone, N., Emanuel, K., Hawcroft, M. K., Hodges, K. I., & Jones, A. (2017). Impacts of hemispheric solar geoengineering on tropical cyclone frequency. *Nature Communications*, 8, 1382.
<https://doi.org/10.1038/s41467-017-01606-0>

Kain, J. S. (n.d.). The Kain–Fritsch Convective Parameterization: An Update. *Journal of Applied Meteorology*, 43(1), 170–181.
[https://doi.org/10.1175/1520-0450\(2004\)043<0170:TKCPAU>2.0.CO;2](https://doi.org/10.1175/1520-0450(2004)043<0170:TKCPAU>2.0.CO;2)

King, R. O., (2013). Financing natural catastrophe exposure: Issues and options for improving risk transfer markets. Congressional Research Service, Library of Congress, 30 pp.

Knapp, K. R., Kruk, M. C., Levinson, D. H., Diamond, H. J., & Neumann, C. J. (2010). The International Best Track Archive for Climate Stewardship (IBTrACS):

Unifying Tropical Cyclone Data. *Bulletin of the American Meteorological Society*, 91(3), 363–376. <https://doi.org/10.1175/2009BAMS2755.1>

Knutson, T. R., McBride, J. L., Chan, J., Emanuel, K., Holland, G., Landsea, C., Held, I., Kossin, J. P., Srivastava, A. K., & Sugi, M. (2010). Tropical cyclones and climate change. *Nature Geoscience*, 3(3), 157–163. <https://doi.org/10.1038/ngeo779>

Knutson, T., Sirutis, J., Garner, S., Vecchi, G., & Held, I. (2008). Erratum: Simulated reduction in Atlantic hurricane frequency under twenty-first-century warming conditions. *Nature Geoscience*, 1(7), 479. <https://doi.org/10.1038/ngeo229>

Korty, R. L., Camargo, S. J., & Galewsky, J. (2012). Tropical Cyclone Genesis Factors in Simulations of the Last Glacial Maximum. *Journal of Climate*, 25(12), 4348–4365. <https://doi.org/10.1175/JCLI-D-11-00517.1>

Korty, R. L., Emanuel, K. A., Huber, M., & Zamora, R. A. (2017). Tropical Cyclones Downscaled from Simulations with Very High Carbon Dioxide Levels. *Journal of Climate*, 30(2), 649–667. <https://doi.org/10.1175/JCLI-D-16-0256.1>

Kossin, J. P., & Camargo, S. J. (2009). Hurricane track variability and secular potential intensity trends. *Climatic Change*, 97(1), 329–337. <https://doi.org/10.1007/s10584-009-9748-2>

Kossin, J. P., Knapp, K. R., Olander, T. L., Velden, C. S. (2020). Global increase in major tropical cyclone exceedance probability over the past four decades.

Proceedings of the National Academy of Sciences, 117 (22) 11975-11980; DOI:
10.1073/pnas.1920849117

Kozar, M. E., Mann, M. E., Emanuel, K. A., & Evans, J. L. (2013). Long-term variations of North Atlantic tropical cyclone activity downscaled from a coupled model simulation of the last millennium. *Journal of Geophysical Research: Atmospheres*, 118(24), 13,313-383,392. <https://doi.org/https://doi.org/10.1002/2013JD020380>

Kravitz, B., MacMartin, D. G., Robock, A., Rasch, P. J., Ricke, K. L., Cole, J. N. S., Curry, C. L., Irvine, P. J., Ji, D., Keith, D. W., Kristjánsson, J. E., Moore, J. C., Muri, H., Singh, B., Tilmes, S., Watanabe, S., Yang, S., & Yoon, J.-H. (2014). A multi-model assessment of regional climate disparities caused by solar geoengineering. *Environmental Research Letters*, 9(7), 74013. <https://doi.org/10.1088/1748-9326/9/7/074013>

Lacis, A. A. and Mishchenko, M. I. (1995), Climate forcing, climate sensitivity, and climate response: A radiative modelling perspective on atmospheric aerosols, in *Aerosol Forcing of Climate* (Charlson R. J. and Heinztenberg, J., (eds.), 416 pp, Wiley, Chichester, pp. 11–42

Liu, K., Lu, H., & Shen, C. (2008). A 1200-year proxy record of hurricanes and fires from the Gulf of Mexico coast: Testing the hypothesis of hurricane–fire interactions. *Quaternary Research*, 69(1), 29–41. <https://doi.org/https://doi.org/10.1016/j.yqres.2007.10.011>

- Lui, Y.S., Tse, L.K.S., Tam, C.Y. *et al* (2021). Performance of MPAS-A and WRF in predicting and simulating western North Pacific tropical cyclone tracks and intensities. *Theoretical Applied Climatology* 143, 505–520.
<https://doi.org/10.1007/s00704-020-03444-5>
- MacMartin, D. G., Wang, W., Kravitz, B., Tilmes, S., Richter, J. H., & Mills, M. J. (2019). Timescale for Detecting the Climate Response to Stratospheric Aerosol Geoengineering. *Journal of Geophysical Research: Atmospheres*, 124(3), 1233–1247. <https://doi.org/https://doi.org/10.1029/2018JD028906>
- Mann, M. E., Woodruff, J. D., Donnelly, J. P., & Zhang, Z. (2009). Atlantic hurricanes and climate over the past 1,500 years. *Nature*, 460(7257), 880–883.
<https://doi.org/10.1038/nature08219>
- Mass, C. F., & Portman, D. A. (1989). Major Volcanic Eruptions and Climate: A Critical Evaluation. *Journal of Climate*, 2(6), 566–593.
[https://doi.org/10.1175/1520-0442\(1989\)002<0566:MVEACA>2.0.CO;2](https://doi.org/10.1175/1520-0442(1989)002<0566:MVEACA>2.0.CO;2)
- Msadek, R., Vecchi, G. A., & Knutson, T. R. (2013). North Atlantic Hurricane Activity: Past, Present and Future. In *Climate Change: Multidecadal and Beyond: Vol. Volume 6* (pp. 285–301). WORLD SCIENTIFIC.
https://doi.org/doi:10.1142/9789814579933_0018
- National Academies of Sciences, Engineering, and Medicine, E. (2019). *Negative Emissions Technologies and Reliable Sequestration: A Research Agenda*. The National Academies Press. <https://doi.org/10.17226/25259>

National Academies of Sciences, Engineering, and M. (2021). *Reflecting Sunlight: Recommendations for Solar Geoengineering Research and Research Governance*.

<https://doi.org/10.17226/25762>

National Research Council. (2015). Climate intervention: Reflecting sunlight to cool earth. In *Climate Intervention: Reflecting Sunlight to Cool Earth*.

<https://doi.org/10.17226/18988>

Oliva, F., Viau, A. E., Peros, M. C., & Bouchard, M. (2018). Paleotemperature database for the western North Atlantic basin. *The Holocene*, 28(10), 1664–1671.

<https://doi.org/10.1177/0959683618782598>

Otto-Bliesner, B. L., Brady, E. C., Fasullo, J., Jahn, A., Landrum, L., Stevenson, S., Rosenbloom, N., Mai, A., & Strand, G. (2016). Climate Variability and Change since 850 CE: An Ensemble Approach with the Community Earth System Model. *Bulletin of the American Meteorological Society*, 97(5), 735–754.

<https://doi.org/10.1175/BAMS-D-14-00233.1>

Paulson, C. A. (1970). The Mathematical Representation of Wind Speed and Temperature Profiles in the Unstable Atmospheric Surface Layer. *Journal of Applied Meteorology and Climatology*, 9(6), 857–861.

[https://doi.org/10.1175/1520-0450\(1970\)009](https://doi.org/10.1175/1520-0450(1970)009)

Pausata, F. S., & Camargo, S. J. (2019). Tropical cyclone activity affected by volcanically induced ITCZ shifts. *Proceedings of the National Academy of Sciences*, 116(16), 7732–7737.

- Pollard, R. T., Rhines, P. B., & Thompson, R. O. R. Y. (1973). The deepening of the wind-Mixed layer. *Geophysical Fluid Dynamics*, 4(4), 381–404.
<https://doi.org/10.1080/03091927208236105>
- Raible, C. C., Brönnimann, S., Auchmann, R., Brohan, P., Frölicher, T. L., Graf, H.-F., Jones, P., Luterbacher, J., Muthers, S., Neukom, R., Robock, A., Self, S., Sudrajat, A., Timmreck, C., & Wegmann, M. (2016). Tambora 1815 as a test case for high impact volcanic eruptions: Earth system effects. *WIREs Climate Change*, 7(4), 569–589. <https://doi.org/https://doi.org/10.1002/wcc.407>
- Rasch, P. J., Crutzen, P. J., & Coleman, D. B. (2008). Exploring the geoengineering of climate using stratospheric sulfate aerosols: The role of particle size. *Geophysical Research Letters*, 35(2). <https://doi.org/https://doi.org/10.1029/2007GL032179>
- Robock, A. (2000). Volcanic eruptions and climate. *Reviews of Geophysics*, 38(2), 191–219. <https://doi.org/https://doi.org/10.1029/1998RG000054>
- Robock, A. (2015). Climate and Climate Change: Volcanoes: Role in Climate. In *Encyclopedia of Atmospheric Sciences: Second Edition* (pp. 105-111). Elsevier Inc..
- S-Y Simon Wang et al (2018). Quantitative attribution of climate effects on Hurricane Harvey's extreme rainfall in Texas. *Environmental Research Letters* 13 054014.
<https://doi.org/10.1088/1748-9326/aabb85>

- Schenkel, B. A., & Hart, R. E. (2011). An Examination of Tropical Cyclone Position, Intensity, and Intensity Life Cycle within Atmospheric Reanalysis Datasets. *Journal of Climate*, 25(10), 3453–3475. <https://doi.org/10.1175/2011JCLI4208.1>
- Schmidt, G. A., Jungclaus, J. H., Ammann, C. M., Bard, E., Braconnot, P., Crowley, T. J., Delaygue, G., Joos, F., Krivova, N. A., Muscheler, R., Otto-Bliesner, B. L., Pongratz, J., Shindell, D. T., Solanki, S. K., Steinhilber, F., & Vieira, L. E. A. (2012). Climate forcing reconstructions for use in PMIP simulations of the Last Millennium (v1.1). *Geoscientific Model Development*, 5(1), 185–191. <https://doi.org/10.5194/gmd-5-185-2012>
- Schurer, A. P., Hegerl, G. C., Mann, M. E., Tett, S. F. B., & Phipps, S. J. (2013). Separating Forced from Chaotic Climate Variability over the Past Millennium. *Journal of Climate*, 26(18), 6954–6973. <https://doi.org/10.1175/JCLI-D-12-00826.1>
- Schurer, A. P., Tett, S. F. B., & Hegerl, G. C. (2014). Small influence of solar variability on climate over the past millennium. *Nature Geoscience*, 7(2), 104–108. <https://doi.org/10.1038/ngeo2040>
- Skamarock, W. C., Klemp, J. B., Dudhia, J., Gill, D. O., Barker, D. M., Duda, M. G., Huang, X.-Y., Wang, W., & Powers, J. G. (2008). *A Description of the Advanced Research WRF Version 3*. <https://doi.org/10.5065/D68S4MVH>
- Stevenson, S., Fasullo, J. T., Otto-Bliesner, B. L., Tomas, R. A., & Gao, C. (2017). Role of eruption season in reconciling model and proxy responses to tropical volcanism.

Proceedings of the National Academy of Sciences, 114(8), 1822 LP – 1826.

<https://doi.org/10.1073/pnas.1612505114>

Strauss, B.H., Orton, P.M., Bittermann, K. et al (2021). Economic damages from Hurricane Sandy attributable to sea level rise caused by anthropogenic climate change. *Nature Communications* 12, 2720.

<https://doi.org/10.1038/s41467-021-22838-1>

Swingedouw, D., Mignot, J., Ortega, P., Khodri, M., Menegoz, M., Cassou, C., & Hanquiez, V. (2017). Impact of explosive volcanic eruptions on the main climate variability modes. *Global and Planetary Change*, 150, 24–45.

<https://doi.org/https://doi.org/10.1016/j.gloplacha.2017.01.006>

Tang, B., & Camargo, S. J. (2014). Environmental control of tropical cyclones in CMIP5: A ventilation perspective. *Journal of Advances in Modeling Earth Systems*, 6(1), 115–128. <https://doi.org/https://doi.org/10.1002/2013MS000294>

Tewari, M., Chen, F., Wang, W., Dudhia, J., LeMone, M. A., Mitchell, K., Ek, M., Gayno, G., Wegiel, J., & Cuenca, R. H. (2004). Implementation and verification of the unified Noah land surface model in the WRF model. *20th Conference on Weather Analysis and Forecasting/16th Conference on Numerical Weather Prediction*.

Ting, M., Camargo, S. J., Li, C., & Kushnir, Y. (n.d.). Natural and Forced North Atlantic Hurricane Potential Intensity Change in CMIP5 Models. *Journal of Climate*, 28(10), 3926–3942. <https://doi.org/10.1175/JCLI-D-14-00520.1>

- United Nations Environment Program. (2017). *The Emissions Gap Report 2017*.
- Vecchi, G. A., & Soden, B. J. (2007). Effect of remote sea surface temperature change on tropical cyclone potential intensity. *Nature*, *450*(7172), 1066–1070.
<https://doi.org/10.1038/nature06423>
- Vecchi, G. A., Landsea, C., Zhang, W., Villarini, G., & Knutson, T. (2021). Changes in Atlantic major hurricane frequency since the late-19th century. *Nature Communications*, *12*(1), 4054.
<https://doi.org/10.1038/s41467-021-24268-5>
- Villarini, G., & Vecchi, G. A. (2013). Projected Increases in North Atlantic Tropical Cyclone Intensity from CMIP5 Models. *Journal of Climate*, *26*(10), 3231–3240.
<https://doi.org/10.1175/JCLI-D-12-00441.1>
- Walsh, K. J. E., Camargo, S. J., Vecchi, G. A., Daloz, A. S., Elsner, J., Emanuel, K., Horn, M., Lim, Y.-K., Roberts, M., Patricola, C., Scoccimarro, E., Sobel, A. H., Strazzo, S., Villarini, G., Wehner, M., Zhao, M., Kossin, J. P., LaRow, T., Oouchi, K., ... Henderson, N. (2015). Hurricanes and Climate: The U.S. CLIVAR Working Group on Hurricanes. *Bulletin of the American Meteorological Society*, *96*(6), 997–1017. <https://doi.org/10.1175/BAMS-D-13-00242.1>
- Wang, Q., Moore, J. C., & Ji, D. (2018). A statistical examination of the effects of stratospheric sulfate geoengineering on tropical storm genesis. *Atmospheric Chemistry and Physics*, *18*(13), 9173–9188.
<https://doi.org/10.5194/acp-18-9173-2018>

- Webb, E. K. (1970). Profile relationships: The log-linear range, and extension to strong stability. *Quarterly Journal of the Royal Meteorological Society*, 96(407), 67–90.
<https://doi.org/https://doi.org/10.1002/qj.49709640708>
- Wehner, M., Prabhat, Reed, K. A., Stone, D., Collins, W. D., & Bacmeister, J. (2015). Resolution Dependence of Future Tropical Cyclone Projections of CAM5.1 in the U.S. CLIVAR Hurricane Working Group Idealized Configurations. *Journal of Climate*, 28(10), 3905–3925. <https://doi.org/10.1175/JCLI-D-14-00311.1>
- Wilson, J. C. et al. (1993). In-situ observations of aerosol and chlorine monoxide after the 1991 eruption of Mount Pinatubo: Effect of reactions on sulfate aerosol, *Science* 261, 1140–1143
- Yan, Q., Korty, R., & Zhang, Z. (2015). Tropical Cyclone Genesis Factors in a Simulation of the Last Two Millennia: Results from the Community Earth System Model. *Journal of Climate*, 28(18), 7182–7202.
<https://doi.org/10.1175/JCLI-D-15-0054.1>
- Yan, Q., Zhang, Z., & Wang, H. (2018). Divergent responses of tropical cyclone genesis factors to strong volcanic eruptions at different latitudes. *Climate Dynamics*, 50(5), 2121–2136. <https://doi.org/10.1007/s00382-017-3739-1>
- Yang, Y., Smith, S. J., Wang, H., Mills, C. M., and Rasch, P. J.: Variability, timescales, and nonlinearity in climate responses to black carbon emissions, *Atmos. Chem. Phys.*, 19, 2405–2420, <https://doi.org/10.5194/acp-19-2405-2019>, 2019.

Zhao, M., Held, I. M., Lin, S.-J., & Vecchi, G. A. (2009). Simulations of Global Hurricane Climatology, Interannual Variability, and Response to Global Warming Using a 50-km Resolution GCM. *Journal of Climate*, 22(24), 6653–6678.
<https://doi.org/10.1175/2009JCLI3049.1>

TABLES

Physics Scheme	Name	Parameter	Value
(1) Micro-physics	WSM6	mp_physics	6
(2) PBL	YSU	bl_pbl_physics	1
(3) Convection	Kain-Fritsch	cu_physics	1
(4) Longwave radiation	RRTMG	ra_lw_physics	4
(5) Shortwave radiation	RRTMG	ra_sw_physics	4
(6) Land surface	Noah	sf_surface_physics	2
(7) Surface Layer	MM5 Similarity	sf_sfclay_physics	91
(8) Ocean	Mixed-layer	sf_ocean_physics	1

Table 1. WRF physics parameterizations used for downscaling. (1) WRF single-moment 6-class (WSM6) for micro-physics (Hong and Lim, 2006), (2) Yonsei University (YSU) for PBL (Hong et al., 2006), (3) Kain-Fritsch for convection (Kain, 2004), (4,5) rapid radiative transfer model with greenhouse gases (RRTMG) for long-wave and short-wave radiation (Iacono et al., 2008), (6) Noah for land surface (Tewari et al., 2002), (7) fifth generation mesoscale model for surface layer (Paulson, 1970; Dyer and Hicks, 1970; Webb, 1970), and (8) simple mixed-layer for ocean (Pollard et al., 1972).

Averages	ERA-I	IBTrACS	Percents	ERA-I	IBTrACS
month	7.65	8.24	May-Nov	0.88	0.99
yearly num	31.27	34.82	0-25 N	0.78	0.73
lat	18.23	21.39	100-50 W	0.65	0.44
lon	-80.72	-88.38	0-40 m/s	0.97	0.70
max wind (m/s)	30.26	34.87	1020-980 hPa	0.81	0.62
min pressure (hPa)	988.56	979.65	(w) 0-100 hrs	0.94	0.91
w-life (hrs)	45.37	44.69	(p) 0-100 hrs	0.94	0.87
p-life (hrs)	44.02	52.53			

Table 2. ERA-I vs. IBTrACS statistics for certain variables. For the first three columns: month indicates the average time of year for hurricanes (for example, the average hurricane occurred slightly after mid-July for ERA-I), yearly num is the average number of hurricanes, lat is the average latitude of hurricanes, lon is the average longitude, max wind is the average maximum wind speed (m/s), min pressure is the average minimum pressure (hPa), w-life is the average time in hours from maximum wind speed to average wind speed of a hurricane (represents decay time), p-life is the same as w-life except for minimum pressure to average pressure of the storm. For the second three columns: May-Nov indicates the percentage of TCs that occur between May and November, 0-25N is the percent of storms that track through this latitude range, 100-50W is the percent of storms that track through this range of longitude, 0-40 m/s is the percent of storms that are in this wind intensity range, 1020-980

hPa indicates the percent of storms in this pressure range, and (w) and (p) 0-100 hrs are the percent of storms that decay within 100 hours.

	All Eruptions D-Value	All Eruptions P-Value	All Eruptions Significance
month	0.0	1.0	0.25
yearly num	0.016	1.0	0.25
lat	0.01	1.0	0.25
lon	0.002	1.0	0.25
max wind (m/s)	0.01	1.0	0.25
min pressure (hPa)	0.006	1.0	0.25
w-life (hrs)	0.004	1.0	0.25
p-life (hrs)	0.0	1.0	0.25
PDI (m ³ /s ²)	0.01	1.0	0.2
ACE (m ² /s ²)	0.01	1.0	0.25

Table 3. KS-tests show a maximum difference between the two samples (D-value) and a probability that the two samples are drawn from the same distribution. Anderson-Darling tests give the level (Significance) at which the null hypothesis can be rejected. The maximum level given by the Anderson-Darling tests is 0.25. (Both Hemispheres)

	All Eruptions D-Value	All Eruptions P-Value	All Eruptions Significance
--	--------------------------	--------------------------	-------------------------------

month	0.0	1.0	0.25
yearly num	0.016	1.0	0.0425
lat	0.014	1.0	0.25
lon	0.016	1.0	0.25
max wind (m/s)	0.02	1.0	0.128
min pressure (hPa)	0.026	1.0	0.068
w-life (hrs)	0.008	1.0	0.25
p-life (hrs)	0.008	1.0	0.25
PDI (m ³ /s ²)	0.026	1.0	0.0175
ACE (m ² /s ²)	0.016	1.0	0.027

Table 4. KS-tests (D- and P-values) and Anderson-Darling tests (Significance) for all eruptions. (Northern Hemisphere)

	5 Strongest Eruptions D-Value	5 Strongest Eruptions P-Value	5 Eruptions Significance
month	0.004	1.0	0.25
yearly num	0.048	0.6	0.001
lat	0.05	0.55	0.001
lon	0.048	0.6	0.001
max wind (m/s)	0.036	0.9	0.001

min pressure (hPa)	0.062	0.28	0.001
w-life (hrs)	0.026	1.0	0.0069
p-life (hrs)	0.024	1.0	0.0125
PDI (m ³ /s ²)	0.05	0.55	0.001
ACE (m ² /s ²)	0.044	0.71	0.001

Table 5. KS-tests (D- and P-values) and Anderson-Darling tests (Significance) for 5 strongest eruptions. (Northern Hemisphere)

	All Eruptions D-Value	All Eruptions P-Value	All Eruptions Significance
month	0.002	1.0	0.25
yearly num	0.01	1.0	0.0059
lat	0.022	1.0	0.119
lon	0.01	1.0	0.25
max wind (m/s)	0.016	1.0	0.25
min pressure (hPa)	0.026	1.0	0.068
w-life (hrs)	0.008	1.0	0.25
p-life (hrs)	0.006	1.0	0.25
PDI (m ³ /s ²)	0.024	1.0	0.03

ACE (m^2/s^2)	0.022	1.0	0.015
-------------------	-------	-----	-------

Table 6. KS-tests (D- and P-values) and Anderson-Darling tests (Significance) for all eruptions. (Southern Hemisphere)

	5 Strongest Eruptions D-Value	5 Strongest Eruptions P-Value	5 Strongest Eruptions Significance
month	0.006	1.0	0.25
yearly num	0.048	0.6	0.001
lat	0.05	0.55	0.001
lon	0.044	0.71	0.001
max wind (m/s)	0.034	0.93	0.00129
min pressure (hPa)	0.06	0.32	0.001
w-life (hrs)	0.026	1.0	0.0069
p-life (hrs)	0.02	1.0	0.055
PDI (m^3/s^2)	0.052	0.5	0.001
ACE (m^2/s^2)	0.042	0.76	0.001

Table 7. KS-tests (D- and P-values) and Anderson-Darling tests (Significance) for 5 strongest eruptions. (Southern Hemisphere)

	All Eruptions
month	0.513
yearly num	0.435
lat	0.494
lon	0.435
max wind (m/s)	0.532
min pressure (hPa)	0.494
w-life (hrs)	0.565
p-life (hrs)	0.506
PDI (m ³ /s ²)	0.552
ACE (m ² /s ²)	0.558

Table 8. Fractional significance tests for all eruptions. The values indicate the fraction of TCs in LME_{control} which exceed the LME_{forced} mean, for a given variable. (Both Hemispheres)

	All Eruptions	10 Strongest Eruptions	5 Strongest Eruptions	1258 Eruption
month	0.597	0.552	0.565	0.623
yearly num	0.539	0.539	0.435	0.753
lat	0.584	0.584	0.786	0.818

lon	0.494	0.435	0.708	0.656
max wind (m/s)	0.662	0.721	0.766	0.877
min pressure (hPa)	0.357	0.383	0.364	0.175
w-life (hrs)	0.662	0.675	0.721	0.63
p-life (hrs)	0.63	0.63	0.747	0.766
PDI (m ³ /s ²)	0.656	0.714	0.799	0.89
ACE (m ² /s ²)	0.656	0.721	0.818	0.935

Table 9. Fractional significance tests for all eruptions, the 10 strongest eruptions, the 5 largest eruptions, and the largest eruption (1258). The values indicate the fraction of TCs in $LME_{control}$ which exceed the LME_{forced} mean, for a given variable. (Northern Hemisphere)

	All Eruptions	10 Strongest Eruptions	5 Strongest Eruptions	1452 Eruption
month	0.422	0.292	0.377	0.156
yearly num	0.39	0.435	0.325	0.169
lat	0.403	0.338	0.396	0.649
lon	0.364	0.318	0.526	0.643
max wind (m/s)	0.429	0.416	0.331	0.461
min pressure (hPa)	0.643	0.617	0.649	0.675

w-life (hrs)	0.422	0.409	0.318	0.305
p-life (hrs)	0.377	0.409	0.37	0.357
PDI (m ³ /s ²)	0.448	0.565	0.617	0.669
ACE (m ² /s ²)	0.455	0.578	0.61	0.721

Table 10. Fractional significance tests for all eruptions, the 10 strongest eruptions, the 5 largest eruptions, and the largest eruption (1452). The values indicate the fraction of TCs in LME_{control} which exceed the LME_{forced} mean, for a given variable. (Southern Hemisphere)

	All Eruptions	90% Confidence Interval	85% Confidence Interval	80% Confidence Interval
month	-0.129	-0.3747,0.1337	-0.3455, 0.1005	-0.323, 0.0754
yearly num	-0.2088	-0.4431,0.0522	-0.4157, 0.0187	-0.3946, -0.0066
lat	0.0176	-0.2417,0.2746	-0.2098, 0.2432	-0.1855, 0.2193
lon	0.1959	-0.0657,0.4323	-0.0321, 0.4045	-0.0068, 0.3832
max wind (m/s)	-0.3176	-0.5322,-0.0647	0.5077, -0.0981	-0.4886, -0.1231
min pressure (hPa)	0.2951	0.0399,0.5142	0.0734, 0.489	0.0985, 0.4695
w-life (hrs)	-0.0875	-0.3381,0.1747	-0.308, 0.1419	-0.2849, 0.117

p-life (hrs)	-0.2927	-0.5122, -0.0373	-0.487, -0.0708	0.4674, -0.0959
PDI (m ³ /s ²)	-0.3577	-0.5633,-0.1104	-0.5405,-0.1427	-0.5223,-0.1674
ACE (m ² /s ²)	-0.3876	-0.5865,-0.1445	-0.5646,-0.1765	-0.5471,-0.2009

Table 11. Correlations between eruption strength and diagnostics for all eruptions in both hemispheres.

	All Eruptions	90% Confidence Interval	85% Confidence Interval	80% Confidence Interval
month	-0.4093	-0.6655, -0.0668	-0.6392, -0.1123	0.6178, -0.147
yearly num	-0.3267	-0.6088, 0.0287	-0.5791, -0.0171	-0.5551, -0.0524
lat	0.0669	-0.2921, 0.4094	-0.2496, 0.3705	-0.2162, 0.3396
lon	0.2673	-0.0936, 0.5661	-0.048, 0.5342	-0.0127, 0.5084
max wind (m/s)	-0.293	-0.5848, 0.0659	-0.5538, 0.0201	-0.5288, -0.0152
min pressure (hPa)	0.2447	-0.1175, 0.5495	-0.0721, 0.5167	-0.0369, 0.4903
w-life (hrs)	-0.0942	-0.432, 0.2667	-0.394, 0.2237	-0.3637, 0.1899

p-life (hrs)	-0.3255	-0.6079, 0.03	-0.5782, -0.0158	-0.5542, -0.0511
PDI (m ³ /s ²)	-0.3592	-0.6314,-0.0081	-0.6031,-0.0539	-0.5801,-0.0891
ACE (m ² /s ²)	-0.4239	-0.6752,-0.0844	-0.6495,-0.1297	-0.6286,-0.1643

Table 12. Correlations between eruption strength and diagnostics for all eruptions in the Northern Hemisphere.

	All Eruptions	90% Confidence Interval	85% Confidence Interval	80% Confidence Interval
month	0.4059	0.0194, 0.6868	0.0706, 0.6588	0.1097, 0.6359
yearly num	0.0702	-0.3283, 0.4475	-0.2818, 0.4056	-0.2451, 0.3721
lat	-0.0491	-0.4304, 0.3471	-0.3877, 0.3012	-0.3537, 0.2649
lon	0.095	-0.3059, 0.4672	-0.2587, 0.4262	-0.2215, 0.3934
max wind (m/s)	-0.3469	-0.6488, 0.0493	-0.6181, -0.0019	-0.5931, -0.0414
min pressure (hPa)	0.3676	-0.0256, 0.6623	0.0256, 0.6325	0.0651, 0.6083
w-life (hrs)	0.021	-0.3716, 0.4072	-0.3266, 0.3636	-0.2909, 0.3288
p-life (hrs)	-0.1701	-0.5249, 0.235	-0.4867, 0.186	-0.456, 0.1476
PDI (m ³ /s ²)	-0.3782	-0.6692,0.0133	-0.6399,-0.0379	-0.616,-0.0773
ACE (m ² /s ²)	-0.3546	-0.6538,0.0405	-0.6235,-0.0107	-0.5987,-0.0502

Table 13. Correlations between eruption strength and diagnostics for all eruptions in the Southern Hemisphere.

FIGURES




Figure 1. Aerosol mass signals for volcanic eruptions from 1150-1850 CE in the Northern Hemisphere. The minimum, 10th largest, and 5th largest signals are shown with dashed lines. All signals were selected as “all NH eruptions” years for this study.

Volcanic Eruptions (SH)




Figure 2. Aerosol mass signals for volcanic eruptions from 1150-1850 CE in the Southern Hemisphere. The minimum, 10th largest, and 5th largest signals are shown with dashed lines. All signals were selected as “all SH eruptions” years for this study.




Figure 3. $\text{LME}_{\text{control}}$ SST power spectrum. 100 years of control data is sufficient in capturing low frequency content.




Figure 4. Downscaled ERA-I hurricane tracks using TSTORMS (A) compared to the IBTrACS database (B) for 1995 through 2005.




Figure 5. Downscaled LME_{forced} hurricane storm tracks using TSTORMS (A) compared to downscaled LME_{control} using TSTORMS (B).




Figure 6. Average number of storms by time of year (month), year, latitude, longitude, average wind speed, average pressure, lifetime based on decay from maximum wind to average wind, lifetime based on decay from minimum pressure to average pressure, accumulated cyclone energy (ACE), and power dissipation index (PDI), between downscaled ERA-I and IBTrACS for 1995 through 2005. Bin widths are 1° , 3° , 1m/s, 1mb for latitude, 6hr, and 6hr for longitude, wind speed, pressure, wind life, and pressure life respectively.

Distributions are normalized by the total number of storms.




Figure 7. Same as Figure 6 except for a comparison between LME_{forced} and $LME_{control}$ for all eruptions in both hemispheres.




Figure 8. Same as Figure 6 except for a comparison between LME_{forced} and $LME_{control}$ for all eruptions in the Northern Hemisphere.




Figure 9. Same as Figure 6 except for a comparison between LME_{forced} and $LME_{control}$ for all eruptions in the Southern Hemisphere.




Figure 10. Same as Figure 6 except for only the five strongest eruptions in the Northern Hemisphere.




Figure 11. Same as Figure 6 except for only the five strongest eruptions in the Southern Hemisphere.